

ADA034156

OFFICE OF NAVAL RESEARCH  
Contract N00014-67-A-0280-0003

Sponsored by the  
Advanced Research Projects Agency  
ARPA Order No. 2721

TECHNICAL REPORT NO. 3

THE TEMPERATURE DEPENDENCE OF TRIPLET STATE REACTION RATE AND  
QUANTUM YIELD FOR AN INTRAMOLECULAR ENONE PHOTOCYCLOADDITION


by

G. Jones, II and B.R. Ramachandran

Prepared for publication  
in the  
Journal of Photochemistry

Department of Chemistry  
Boston University  
Boston, Massachusetts 02215

December 1, 1976



Reproduction in whole or in part is permitted for any purpose  
of the United States Government

Approved for public release; distribution unlimited

The views and conclusions contained in this document are those of the authors  
and should not be interpreted as necessarily representing the official policies,  
either expressed or implied, of the Advanced Research Projects Agency of the  
U.S. Government.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
Technical Report No. 3	1 May - 31 Oct 75	
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
The Temperature Dependence of Triplet State Reaction Rate and Quantum Yield for an Intramolecular Enone Photocycloaddition		Technical, 5/1/75-10/31/75
7. AUTHOR(s)		6. PERFORMING ORG. REPORT NUMBER
G. Jones, II and B.R. Ramachandran		
9. PERFORMING ORGANIZATION NAME AND ADDRESS		8. CONTRACT OR GRANT NUMBER(s)
Department of Chemistry, Boston University Boston, Massachusetts 02215		NO 0014-67-A0280-0003 ARPA Order - 2722
11. CONTROLLING OFFICE NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217		ARPA Order No. 2721/1-4-74 Program Code No. 4F10
12. REPORT DATE		13. NUMBER OF PAGES
December 1, 1976		6
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
Guilford Jones, II Bellampalli R. / Ramachandran		unclassified
16. DISTRIBUTION STATEMENT (of this Report)		
Reproduction in whole or in part is permitted for any purpose of the United States Government. Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
Prepared for publication in the Journal of Photochemistry		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Valence photoisomerization, triplet state reaction, temperature dependence of excited state lifetime, diradicals		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
<p>Valence photoisomerization (320-380 nm) of endo-tricyclo(5.2.1.0<sup>2,6</sup>)deca-4,8-diene-3-one has been studied as a function of temperature. At 4 - 51° the isomerization quantum yield is independent of temperature (<math>\Phi = 0.4</math>). In the same range the lifetime for the triplet state responsible for isomerization decreases. Arrhenius parameters for triplet decay are computed, <math>E_a = 4.7\text{kcal/mol}</math>; <math>\log A = 13.7</math>. A mechanism involving formation of a diradical intermediate which partitions to the valence isomers is suggested.</p>		

DD FORM 1473  
1 JAN 73EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

401003



THE TEMPERATURE DEPENDENCE OF TRIPLET STATE REACTION RATE AND QUANTUM YIELD FOR AN INTRAMOLECULAR ENONE PHOTOCYCLOADDITION.

Guilford Jones, II\* and Bellampalli R. Ramachandran

Department of Chemistry, Boston University, Boston, Massachusetts 02215 (U.S.A.)

APPROVED FOR	
NTIS	Microfilm <input checked="" type="checkbox"/>
DPC	Book <input type="checkbox"/>
ORAL REPORT	<input type="checkbox"/>
DISTRIBUTION/AVAILABILITY CODES	
Dist.	Avail. and/or Special
A	

Although photochemically induced rearrangements of ketones have been the focus of numerous studies,<sup>1</sup> few reports have documented a temperature dependence of excited state reaction rates or of quantum yields.<sup>2</sup> We wish to provide Arrhenius parameters for reactive decay of an excited enone. Our study of the kinetics of valence isomerization  $1 \rightarrow 2$  is the first detailed look at an intramolecular version of cyclopentenone photoannulation,<sup>3</sup> and is of additional interest since the  $1 - 2$  couple is a prototype photon energy storage system.<sup>4</sup> We note particularly that the high rate of reaction of triplet  $1$  is moderately temperature dependent while the quantum yield for isomerization is not.

Isomerization  $1 \rightarrow 2$  is clean, quantum-efficient, highly endoergic, moderately photochromic, and reversible upon treatment at elevated temperatures with Rh(I) catalysts.<sup>4</sup> The reaction may be sensitized as shown in Table I. The effectiveness of acetophenone vs. anthrone is consistent with a triplet energy for  $1$ ,  $E_T = 72.9$  kcal/mol, estimated from its emission at 77°K in EPA glass ( $\lambda_{\max} 392$  nm,  $1680$  cm<sup>-1</sup> vibrational spacing).<sup>6</sup> The results also point to a unit quantum efficiency for intersystem crossing which precedes photorearrangement on direct irradiation of  $1$ .

Isomerization was quenched by 2,5-dimethyl-2,4-hexadiene ( $E_T < 58$  kcal/mol), and plots of  $\phi_0/\phi$  vs.  $[Q]$  were linear. The results of Stern-Volmer analysis along with quantum yields and rate constants for diffusion controlled quenching by diene ( $k_q$ ) for different temperatures are shown in Table II.

Assuming a mechanism for photoisomerization in which  $T_1$  partitions with inter-






Table I. Quantum Yields for Sensitized Isomerization  $\lambda \rightarrow \lambda^a$

Sensitizer (M) <sup>b</sup>	E <sub>T</sub> (kcal/mol)	$\phi^c$
---	---	0.37 $\pm$ 0.02
acetophenone (0.3)	73.6	0.33 $\pm$ 0.02
anthrone (0.07)	71.9	0.06 $\pm$ 0.004

<sup>a</sup> Rayonet reaction, 3500 lamps (325 - 385 nm), valerophenone actinometer (ref 5), 0.07 M  $\lambda$  in benzene. <sup>b</sup> > 80% light absorbed by sensitizer. <sup>c</sup> Corrected for competitive and differential absorption.

system crossing directly to  $\lambda$  ( $k_r$ ) and to  $\lambda$  ( $k_d$ ), one applies the relationships,  $\phi_0/\phi = 1 + k_q[Q]/k_r + k_d$  and  $\phi_0 = k_r/k_r + k_d$  and finds that (at 30°),  $k_r = 8.0 \times 10^9 \text{ sec}^{-1}$  and  $k_d = 1.4 \times 10^{10} \text{ sec}^{-1}$ . The calculated  $k_d$  is unacceptable in view of the size of radiationless decay constants determined for other enones ( $k_d = 10^7 - 10^8 \text{ sec}^{-1}$ ).<sup>3a</sup> Treatment of the data using the Arrhenius equation gives the preexponential factors,  $\log A \sim 13$ , for both  $k_r$  and  $k_d$ . These values appear improbably high for intercombinational processes.

In an alternative scheme,  $T_1$  does not lead directly to  $\lambda$  or  $\lambda$  but gives triplet diradial  $\lambda$ ,<sup>7</sup> which partitions to the isomers. The relationship,  $\phi_0/\phi = 1 + k_q[Q]/k_r'$ , then applies, with which  $k_r'$  may be obtained (results shown in Table II). An Arrhenius plot of the rate data is shown in the Figure along with calculated parameters for  $k_r'$ .

For the diradical mechanism, absolute quantum yields for photorearrangement are determined by the relative rates of closure ( $k_i$ ) and cleavage ( $k_c$ ) of  $\lambda$ , and  $(1/\phi_0) - 1 = k_c/k_i$ . A plot of  $(1/\phi_0) - 1$  vs.  $1/T$  shows that  $E_a(c) - E_a(i) = 0.09 \pm 0.17 \text{ kcal/mol}$  and  $A(c) / A(i) = 2.0 \pm 0.6$ .

The quantum yield for photoisomerization is understood then in terms of frequency

Table II. Direct Irradiation Quantum Yields and Quenching Data for Isomerization

$\lambda \rightarrow \lambda'$				
$T(^{\circ}C)^a$	$\phi$	Slope ( $M^{-1}$ ) <sup>b</sup>	$k_q (M^{-1} sec^{-1})^c$ $\times 10^{-9}$	$k_r' (sec^{-1})$ $\times 10^{-10}$
4	---	$0.395 \pm 0.025$	4.15	$1.05 \pm 0.07$
18	$0.38 \pm 0.02$	$0.345 \pm 0.015$	5.38	$1.56 \pm 0.07$
30	$0.37 \pm 0.02$	$0.310 \pm 0.015$	6.72	$2.17 \pm 0.11$
40	$0.36 \pm 0.02$	$0.275 \pm 0.015$	7.84	$2.85 \pm 0.15$
51	$0.38 \pm 0.02$	$0.265 \pm 0.015$	9.33	$3.52 \pm 0.20$

<sup>a</sup>  $\pm 1^{\circ}C$ . <sup>b</sup> From plots of  $\phi_0/\phi$  vs.  $[Q]$ ; irradiation at 325-385 nm in cyclooctane solution. <sup>c</sup> Calculated using the Debye equation and measured viscosities.

factor differences favoring diradical cleavage over closure.<sup>6,7</sup> The diradical mechanism is economical, it leads to an informative treatment of the quantum yield and quenching data, but it is not uniquely consistent with the facts.<sup>8</sup> We will discuss in a full paper the possibilities that the thermal barrier to photoisomerization separates an initial enone triplet from a reactive (unquenchable) upper triplet of different configuration or from a conformation which is equivalent to the exciplex in intermolecular photoannulation<sup>3a</sup> (in which enone and olefin chromophores interact strongly)<sup>10</sup>.

This research was supported by the Advanced Research Projects Agency of the Department of Defense (monitored by ONR) and by the Petroleum Research Fund.






Figure. Arrhenius plot and parameters, indicating the temperature dependence for reactive decay of triplet I ( $k_r'$ ).

- (1) See, for example, (a) S.S. Hixson, P.S. Mariano, and H.E. Zimmerman, Chem. Rev., **73**, 531 (1973); (b) K. Schaffner in "Organic Reactions in Steroid Chemistry," vol 2, Ed., J. Fried and J.A. Edwards, Van Nostrand Reinhold Co., N.Y., 1972, p 288; (c) H.E. Zimmerman, Angew. Chem. Intern. Ed. Engl., **8**, 1 (1969).
- (2) (a) H.E. Zimmerman and W.R. Elser, J. Amer. Chem. Soc., **91**, 887 (1969); (b) C.D. DeBoer, W.G. Herkstroeter, A.P. Marchetti, A.G. Schultz, and R.H. Schlessinger, ibid., **95**, 3963 (1973)
- (3) (a) P. de Mayo, Accounts Chem. Res., **4**, 41, (1971); (b) P.E. Eaton, ibid., **1**, 50 (1968); (c) P.G. Bauslaugh, Synthesis, **2**, 287 (1970)
- (4) G. Jones, II and B.R. Ramachandran, J. Org. Chem., in press.
- (5) P.J. Wagner, P.A. Kelso, and R.G. Zepp, J. Amer. Chem. Soc., **94**, 7480 (1972).
- (6) Sensitization of isomerization 1  $\rightarrow$  2 by the  $T_2$  state of substituted anthracenes ( $E_{T_2} \sim 73 - 74$  kcal/mol) has been reported; see R.O. Campbell and R.S.H. Liu, J. Amer. Chem. Soc., **95**, 6560 (1973), and R.S.H. Liu and J.R. Edman ibid., **91**, 1492 (1969).
- (7) (a) For a summary of mechanistic possibilities for enone photoannulation including the diradical route, see ref 3a. (b) We have chosen for illustration the more stable diradical with no experimental basis for excluding intervention of its regioisomer.
- (8) For intermolecular photoannulation direct formation of diradical from triplet enone (plus olefin), without intervention of an exciplex has been considered doubtful due to the high rates of addition ( $k_r$ 's  $\sim 10^8 - 10^9$  M<sup>-1</sup> sec<sup>-1</sup>) compared to the bimolecular rates of olefin - radical reactions.<sup>3a</sup> The free radical, ground state model may be unsatisfactory however, in view of the likely greater exothermicity of photoaddition and if the enone triplet utilizes a pyramidal<sup>9</sup> reactive center (odd e density in a directed orbital), for attack on olefin. For 1, we also expect an especially high rate of intramolecular triplet enone addition, enhanced by a high frequency factor (by comparison, much more favorable entropy of activation) as observed.
- (9) K. Wiesner, Tetrahedron, **31**, 1655 (1975).
- (10) Although enone and olefin chromophores (according to models) are separated by only 2.5 - 3.0 Å, there is no indication in the uv absorption or low temperature emission spectra of significant interaction in the ground state or lowest triplet equilibrium geometries.